AD-A261 137

ATION PAGE

Form Approved

OMB No. 0704-0188

ng the collection of information. Send commit to Washington Headquarters Services, Direct		nformation. Send comments re dquarters Services, Directorate	e for reviewing instructions, searching existing data sources, is regarding this burden estimate or any other aspect of this rate for information Operations and Reports, 1215 Jefferson on Project (0704-0188), Washington, OC 20503.	
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 12/28/92	3. REPORT TYPE A	ND DATES COVERED	
4. TITLE AND SUBTITLE Mechanisms of Decomp of Borane	osition and Oxid	dation	5. FUNDING NUMBERS DAALO3+88-K-0187	
6. AUTHOR(S) S.H. Baue	r			
7. PERFORMING ORGANIZATION NAME(Cornell University - Baker Chemical Labor	8. PERFORMING ORGANIZATION REPORT NUMBER			
Ithaca, New York 14	853-1301			
9. SPONSORING/MONITORING AGENCY U.S. Army Research Offi P.O. Box 12211)	10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
Research Triangle Park, NC 27709-2211			ARO 25523.2-CH	

11. SUPPLEMENTARY NOTES

The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

126. DISTRIBUTION CODE

EB 19 1993

13. ABSTRACT (Maximum 200 words)

The rates of gas phase reactions between boranes (B2H6; B5H9; B10H14) and oxygen atoms (3P state) were studied (~350K) in a flow-tube reactor by determining time dependent levels of OH species, as functions of reactant concentrations. Quantitative fitting of these data for B2H6 and B5H9 led to mechanisms, and estimated rate constants, for the direct attack on the boranes, and for reactions of O(3 P) with fragmentation products (in particular, HBO and HBOH). A degradation sequence was established for each. Measurements (via LIF) of [OH] from B10H14 were not completed; our data show that under comparable conditions the deca produces ~ 60 fold higher [OH] than does penta. Our QPMS diagnostic of a newly designed flow-tube system and revitalized massspectrometer permitted evaluation of a rate constant for the net destruction of B5H9, and qualitative estimates of relative concentrations of B5H9 fragmentation products.

14. SUBJECT TERMS Borane LIF of product OH Combustion with O	15. NUMBER OF PAGES 2 16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED	UL

STATEMENT OF THE PROBLEM:

- A). To develop general mechanisms for the reactions of the more stable boranes with atomic $O(^3P)$, at ambient temperatures.
- B). To measure quantitatively time dependent OH concentrations, via laser induced fluorescence, generated as the dominant product species, rather than measure overall rates of degradation of the fuels.
- C). To develop fragmentation patterns and estimate rate constants for the dominant reactions in the multi-step mechanisms, comparable to those available for the combustion of low molecular weight hydrocarbons.
- D). To develop a reliable QPMS diagnostic for the effluents from a well designed flow-reactor, for quantitative assaying of reaction intermediates, generated during the combustion process.

SUMMARY OF RESULTS:

- a) A low pressure (5 20 Torr) flow reactor was set up; it provides controlled flow rates of measured levels of oxygen atoms, the fuels of interest, and inert carrier gases.
- b) A Nd/YAG/Dye laser system was set up for generating pulses of 307.93 nm radiation, to excite a Q-branch line of the ground state of OH to the A state, from which fluorescence is emitted.
- c) Fluorescence radiation detection (photomultiplier with filter) digitizing, recording and data processing system was assembled. The photube signal is sent to a BIOMATION 8200, a dual channel Tracor-Northern, and on to a computer for data analysis and plotting.
- d) Appropriate thermochemical and kinetic codes were assembled. The available codes for integrating multiple sets of coupled rate equations were extended and tested.
- e) Numerous preliminary experiments were run with selected hydrocarbons. C2H6 was chosen as a standard for calibrating the output signals, so that the fluorescence intensities recorded for the B/H species could be placed on a quantitative basis, (in mole/liter of OH), that appeared at the sampling pin-hole, centrally located in the flow-tube, and on to the intersection point of the jet with the laser beam. Reaction times were calculated from measured flow rates and location of the fuel injector.
- f) A self-consistent, quantitatively scalable mechanism was developed for oxygen atom attack on ethane.
 - g) A similar mechanism was developed for diborane.
 - h) A similar mechanism was developed for pentaborane(9); it



incorporated the reactions and rate constants used for diborane.

i) Rates of OH production in the reactions of $O(^3P)$ with H3BCO and B10H14 were recorded for limited combinations of fuel concentrations and temperatures. These are insufficient for postulating break-down sequences. However, under comparable conditions, B10H14 produces ≈ 60 fold higher levels of OH than does B5H9. Steady-state [OH] is attained in B10H14 at 18 ms post-mixing, whereas steady-state with B5H9 appears at 4 ms.

PERSONNEL (Post-doctoral associates):

```
Dr. Huy-Zu Cheng {05/02/89 - 05/01/92}
Dr. Julio Caballero {05/14/90 - 07/31/91}
Dr. Cahn Park {08/19/91 - 05/01/92}
Dr. Scott Hurst {06/02/92 - 10/14/92}
```

PUBLICATIONS:

Relaxation Kinetics in Lewis Acid:base Systems --F3B + O(CH3)2 = F3B:O(CH3)2
D.B. Borchard, K. Saito and S.H. Bauer
J. Molecular Structure 223, 375 (1990)

Chemiluminescent Reactions of Boron Hydrides with O(3P) --H-Z. Cheng, P.M. Jeffers and S.H. Bauer
Combustion Institute, Eastern Section Proceedings (Oct. 1991)

O(³P) Attack on Boron Hydrides: B2H6 and B5H9 H-Z. Cheng and S.H. Bauer J. Chem. Physics, 94, 7076 (1991)

On the Exoegicities of Borane Combustion; Thermochemical Potential vs. Kinetic Constraints
S.H. Bauer and H-Z. Cheng
Submitted to: International J. Chem. Kinetics.

DITC QUALLTY DISPECTED 3

Acces	ssion For	
NTIB	A&I	3
D: E		
	গুলিকী	
J i; . :	n_	
, 		
11201	Marchay	
100	`î L. `î %y :	Coárs
.	Section 1739 के,	
Dist	ំ ខំព្ ខែរ	. }
01		j
Ν,	1 1]
•	1 1	3